




*Baileya ophthalmica* Eyed Baileya




*n*=23  
High Mt.



*n*=74  
Low Mt.



*n*=80  
*Pd*




NC counties: 43

High counts of:

- 51 - Montgomery - 2007-08-09
- 43 - Pender - 1998-07-01
- 34 - Rutherford - 2013-05-09

Status	Rank
NC	US
NC	Global



*n*=53  
*CP*

Three periods to each month: 1-10 / 11-20 / 21-31

FAMILY: Nolidae SUBFAMILY: Risobinae TRIBE:  
TAXONOMIC\_COMMENTS:

FIELD GUIDE DESCRIPTIONS: Covell (1984); Beadle and Leckie (2012)

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS:

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS:

DISTRIBUTION: Probably occurs statewide

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT: Our records come primarily from wet-mesic forests but there are also some from drier ridges and summits.

FOOD: Larvae feed on American Hornbeam (<i>Carpinus caroliniana</i>), American Hop-hornbeam (<i>Ostrya virginiana</i>), American Hazelnut (<i>Corylus americana</i>), and Beaked Hazelnut (<i>Corylus cornuta</i>) (Wagner et al., 2011). We do not have any feeding records in North Carolina.

OBSERVATION\_METHODS:

NATURAL HERITAGE PROGRAM RANKS: G5 S4S5

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: