




Hyperstrotia flaviguttata Yellow-spotted Graylet



FAMILY: Erebidae SUBFAMILY: Phytometrinae TRIBE:
TAXONOMIC COMMENTS:

FIELD GUIDE DESCRIPTIONS: Covell (1984); Beadle and Leckie (2012)

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS:

TECHNICAL DESCRIPTION, IMMATURE STAGES: Wagner et al. (2011)

ID COMMENTS:

DISTRIBUTION: Primarily found in the Coastal Plain in North Carolina, but with scattered records from the Piedmont and one from the mountains.

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT: Associated with barrens in the north but becomes more widespread further south (Wagner et al., 2011). Our records come mainly from dry sandridge habitats in the Coastal Plain. We also have one record from a mafic glade in the Piedmont and a couple from dry ridges in the Mountains.

FOOD: Larvae feed on oaks (Wagner et al., 2011).

OBSERVATION METHODS:

NATURAL HERITAGE PROGRAM RANKS: G4 SNR [S3S4]

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: