





Marimatha nigrofimbria Black-bordered Lemon Moth




High counts of:

- 150 - Carteret - 2013-08-21
- 50 - Stokes - 1998-08-18
- 50 - Stokes - 1998-08-18

● = Sighting or Collection in NC since 2003
● = Not seen since 2003

| Status | Rank |
|--------|--------|
| NC | US |
| NC | Global |



FAMILY: Noctuidae SUBFAMILY: Eustotiinae TRIBE:
TAXONOMIC_COMMENTS:

FIELD GUIDE DESCRIPTIONS: Covell (1984; as *Thioptera nigrofimbria*); Beadle and Leckie (2012)

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS:

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS:

DISTRIBUTION: Found statewide

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT:

FOOD: Oligophagous, feeding on grasses. Wagner et al. (2013) specifically mention Crab Grass, with Panic Grass and Bluegrass in captivity. The range of habitats used by this species in the state suggest that a wide variety of grasses are used.

OBSERVATION_METHODS:

NATURAL HERITAGE PROGRAM RANKS: G5 S5

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: