




*Apamea amputatrix* Yellow-headed Cutworm Moth



FAMILY: Noctuidae SUBFAMILY: Noctuinae TRIBE: Apameini

TAXONOMIC COMMENTS: A Holarctic genus of about 140 species, with 63 recorded in North America, mostly from boreal or montane habitats (Mikkola et al., 2009). Seventeen species occur in North Carolina with the majority of records coming from the mountains.

FIELD GUIDE DESCRIPTIONS: Covell (1984); Beadle and Leckie (2012)

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS:

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS:

DISTRIBUTION: Please refer to the dot map.

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT: Our records come mainly from high elevations, where they are associated with high elevation forests, bald-like semi-natural grasslands, and alluvial habitats.

FOOD: Larvae feed on a wide range of herbaceous plants, including both grasses and forbs; a number of crop species are used, including cabbage, lettuce, corn, currants, roses, spinach, and wheat (Mikkola et al., 2009). We do not have any feeding records in North Carolina.

OBSERVATION\_METHODS:

NATURAL HERITAGE PROGRAM RANKS: G5 S3S4

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: