




Xylomoia chagnoni Reed Canary Grass Borer Moth



| Status | Rank | | |
|--------|------|----|--------|
| NC | US | NC | Global |

n=0
CP



Three periods to each month: 1-10 / 11-20 / 21-31

FAMILY: Noctuidae SUBFAMILY: Noctuinae TRIBE: Apameini

TAXONOMIC COMMENTS:

FIELD GUIDE DESCRIPTIONS: Beadle and Leckie (2012)

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS:

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS:

DISTRIBUTION: Please refer to the dot map.

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT: Our two records come from riparian habitats in the Mountains.

FOOD: Larvae feed on Reed Canary-grass (<i>Phalaris arundinacea</i>) (Wagner et al., 2011).

OBSERVATION METHODS:

NATURAL HERITAGE PROGRAM RANKS: GNR SNR [SU]

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: This is a northern species with our records representing what appears to be a disjunct population, possibly reaching our area along the New River/Ohio River drainage. Its host plant, however, is widespread in the Mountains and the moth should be looked for in areas further to the south.