




Globia oblonga Oblong Sedge Borer Moth



Status Rank
NC US NC Global



Three periods to each month: 1-10 / 11-20 / 21-31

FAMILY: Noctuidae SUBFAMILY: Noctuinae TRIBE: Apameini

TAXONOMIC COMMENTS:

FIELD GUIDE DESCRIPTIONS: Covell (1984; as *Archana oblonga*); Beadle and Leckie (2012)

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS:

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS:

DISTRIBUTION: Please refer to the dot map.

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT: Our records come from marshy beaver and depression ponds, lakeshores, mountain bogs, and other open wetlands that support cattails.

FOOD: Larvae reportedly feed on cattails, including Broadleaf Cattail (*Typha latifolia*) and Narrowleaf Cattail (*T. angustifolia*) (Eiseman, 2022).

OBSERVATION METHODS:

NATURAL HERITAGE PROGRAM RANKS: G5 SNR [S2S3]

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: