




# *Amphipoea americana* American Ear Moth



FAMILY: Noctuidae SUBFAMILY: Noctuinae TRIBE: Apameini  
 TAXONOMIC\_COMMENTS:

FIELD GUIDE DESCRIPTIONS: Covell (1984); Beadle and Leckie (2012)

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS:

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS: Very similar in color and markings to *Amphipoea interoceanica*, but is larger and not as stocky (Forbes, 1954).

DISTRIBUTION: Please refer to the dot map.

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT: Our records come mainly from high elevation forests and semi-natural grasslands, but also from wet meadows and bogs. One isolated Piedmont record comes from a residential area adjacent to both fields and bottomlands.

FOOD: Larvae reportedly feed on sedges and grasses, including corn, but have also been recorded on iris (Wagner et al., 2011). We do not have any feeding records in North Carolina.

OBSERVATION\_METHODS:

NATURAL HERITAGE PROGRAM RANKS: G5 S4S5

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: