



Papaipema polymniae Cup Plant Borer Moth



FAMILY: Noctuidae SUBFAMILY: Noctuinae TRIBE: Apameini

TAXONOMIC COMMENTS: One of 44 species in this genus that occur in North America north of Mexico (Lafontaine and Schmidt, 2010, 2015), 30 of which have been recorded in North Carolina

FIELD GUIDE DESCRIPTIONS:

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS:

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS:

DISTRIBUTION: Please refer to the dot map.

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT: Piedmont records come from rich, wet-to-mesic forests. Mountain records come from cove forests and also from a high elevation site that has extensive open fields as well as stands of hardwood forests.

FOOD: Larvae appear to be monophagous on Hairy Leafcup (<i>Smallanthus uvedalia</i>) (Wagner et al., 2011), which has been confirmed as the host in North Carolina.

OBSERVATION METHODS:

NATURAL HERITAGE PROGRAM RANKS: G4? S2S4 [S2S3]

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: