



Papaipema baptisiae Indigo Stem Borer Moth



FAMILY: Noctuidae SUBFAMILY: Noctuinae TRIBE: Apameini

TAXONOMIC COMMENTS: One of 44 species in this genus that occur in North America north of Mexico (Lafontaine and Schmidt, 2010, 2015), 30 of which have been recorded in North Carolina

FIELD GUIDE DESCRIPTIONS: Beadle and Leckie (2012)

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS:

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS:

DISTRIBUTION: Please refer to the dot map.

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT: Our records come from forest-field ecotones found across the state

FOOD: Reported hosts include dogbane (*Apocynum*), wild indigo (*Baptisia*), Mayapple (*Podophyllum peltatum*), dock (*Rumex*), and ironweed (*Vernonia*) (Wagner et al., 2011).

OBSERVATION_METHODS:

NATURAL HERITAGE PROGRAM RANKS: G4 SNR [S3S4]

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: