




# *Dypterygia rozmani* American Bird's-wing Moth



Status	Rank		
NC	US	NC	Global



Three periods to each month: 1-10 / 11-20 / 21-31

FAMILY: Noctuidae SUBFAMILY: Noctuinae TRIBE: Dypterygiini

TAXONOMIC COMMENTS:

FIELD GUIDE DESCRIPTIONS: Covell (1984); Beadle and Leckie (2012)

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS:

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS:

DISTRIBUTION: Please refer to the dot map.

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT: The use of Dock and Plantain suggest that ruderal or other early successional habitats could be used. However, our records come mainly from mesic-to-wet forests, which is more consistent with the use of Common Climbing Buckwheat.

FOOD: Larvae have been reported on dock (<i>Rumex</i> spp.), Common Climbing Buckwheat (<i>Fallopia scandens</i>), and plantain (<i>Plantago</i> spp.) (Robinson et al., 2010; Wagner et al., 2011).

OBSERVATION METHODS:

NATURAL HERITAGE PROGRAM RANKS: G5 SNR [S2S3]

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: