




Crambodes talidiformis Verbena Moth



FAMILY: Noctuidae SUBFAMILY: Condicinae TRIBE: Leuconyctini
 TAXONOMIC_COMMENTS:

FIELD GUIDE DESCRIPTIONS: Covell (1984); Beadle and Leckie (2012)

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS:

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS:

DISTRIBUTION: Please refer to the dot map.

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT: Wagner et al. (2011) list open areas as habitats used by this species, including fields, beaches, waste lots, sand plains, and wetlands. Our records come almost entirely from the Mountains, with most from wet meadows or mountain bogs where Blue Vervain could be the host. A few also come from high elevation mesic forests where White Vervain could be used.

FOOD: Larvae feed on vervains (<i>Verbena</i> spp.) (Wagner et al., 2011). We do not have any feeding records in North Carolina.

OBSERVATION_METHODS:

NATURAL HERITAGE PROGRAM RANKS: G5 SNR [S2S3]

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: