


# *Spodoptera latifascia* Velvet Armyworm Moth




*n=0*  
High Mt.

*n=0*  
Low Mt.

High counts of:


- 1 - Cumberland - 2002-09-18
- 1 - Moore - 2009-06-17
- 1 - Mecklenburg - 2019-05-29



Status Rank  
NC US NC Global

*n=3*  
*Pd*

*n=2*  
*CP*



FAMILY: Noctuidae SUBFAMILY: Noctuinae TRIBE: Prodeniini  
TAXONOMIC\_COMMENTS:

FIELD GUIDE DESCRIPTIONS:

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS:

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS: According to Pogue (2002), *latifascia* can easily be confused with *S. ornithogalli* but the spots of adterminal line are spindle-shaped in *S. latifascia* and crescent-shaped in *S. ornithogalli*. The veins of the hindwing are white in *latifascia* but are shaded with brown in *ornithogalli*.

DISTRIBUTION: Please refer to the dot map.

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT: Wagner et al. (2011) list croplands, waste areas, and other open lands as habitats used by this species. Our records come open wetlands, sand ridges, and residential areas, all of which may represent migrants from the south.

FOOD: Like other members of this genus, larvae of this species are broadly polyphagous (Wagner et al., 2011)

OBSERVATION\_METHODS:

NATURAL HERITAGE PROGRAM RANKS: GNR SNA

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: