




# Condica confederata Confederate Moth



FAMILY: Noctuidae SUBFAMILY: Condicinae TRIBE: Condicini  
 TAXONOMIC\_COMMENTS:

FIELD GUIDE DESCRIPTIONS: Covell (1984); Beadle and Leckie (2012)

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS: Poole (2017)

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS: Males can be distinguished from Condica cupentia by forewing pattern but females are more safely distinguished by genitalic features (Poole, 2017).

DISTRIBUTION: Please refer to the dot map.

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT: The majority of our records come from open herb-dominated habitats in the Coastal Plain, including barrier islands, coastal fringe sandhills, and sandhill seeps. Habitats are mostly dry but have imbedded wetlands or marshes nearby.

FOOD: Larvae feed on Climbing Hempvine (Mikania) (Wagner et al., 2011)

OBSERVATION\_METHODS:

NATURAL HERITAGE PROGRAM RANKS: G5 S3S4

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: