




Lithophane laticinerea Broad Ashen Pinion



Status	Rank		
NC	US	NC	Global



Three periods to each month: 1-10 / 11-20 / 21-31

FAMILY: Noctuidae SUBFAMILY: Noctuinae TRIBE: Xylenini

TAXONOMIC COMMENTS: One of 51 species in this genus that occur in North America (Lafontaine and Schmidt, 2010, 2015), 25 of which have been recorded in North Carolina

FIELD GUIDE DESCRIPTIONS: Beadle and Leckie (2012)

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS:

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS:

DISTRIBUTION: Please refer to the dot map.

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT: Our two records come from wet forests, one from a montane riparian forest and one from a non-riverine swamp forest in the Coastal Plain

FOOD: Larvae feed primarily on maples, including Red Maple (*Acer rubrum*) and Silver Maple (*A. saccharinum*); other confirmed records come from American Basswood (*Tilia americana*), crabapple (*Malus*), and Black Oak (*Quercus velutina*) (Wagner et al., 2011).

OBSERVATION METHODS:

NATURAL HERITAGE PROGRAM RANKS: GNR SNR [S1S2]

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: