


Epiglaea apiata Pointed Sallow



FAMILY: Noctuidae SUBFAMILY: Noctuinae TRIBE: Xylenini

TAXONOMIC COMMENTS: The genus contains two species found principally in the Eastern United States and in North Carolina. The two species are not closely related and *E. decliva* will eventually be moved to another genus.

FIELD GUIDE DESCRIPTIONS: Covell (1984)

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS: Forbes (1954)

TECHNICAL DESCRIPTION, IMMATURE STAGES: Wagner et al. (2011)

ID COMMENTS: Very much like a *Metaxaglaea* species but the vestiture is finer and the lower lobe of the reniform contains a distinct black spot. The subterminal line is reddish in the medial portion and stands out.

DISTRIBUTION: All of our records come from the Outer Coastal Plain and Sandhills.

FLIGHT COMMENT: Univoltine, with records in October and November.

HABITAT: All but one of our records come from Longleaf Pine habitats, including savannas, flatwoods, and sandhills. The one exception comes from an area of extensive peatlands, with no Longleaf Habitat located within several miles of the site. Most of the Longleaf sites where this species has been recorded also have imbedded pocosins, pond pine woodlands, or stands of Atlantic White Cedar, suggesting that *E. apiata* feeds on peatland heaths, as well as those associated with the sandier, more frequently-burned habitats characterized by the presence of Longleaf Pines.

FOOD: Recorded on Cranberry and captive larvae feed on various *Vaccinium* species (Forbes, 1954). Reported to be a pest occasionally on cranberries. Wagner et al. (2011) also report Leatherleaf (*Chamaedaphne calyculata*) and huckleberry (*Gaylussacia*). We do not have any specific hostplant records from North Carolina.

OBSERVATION_METHODS: Readily attracted to light and presumably comes to bait though records are lacking.

NATURAL HERITAGE PROGRAM RANKS: G5 [S3S4]

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: This species appears to be associated with just a narrow range of Longleaf and peatland habitats, all of which have declined drastically since European colonization and which are still threatened in the state by continued habitat conversion and fire suppression. As with other species associated with fire-maintained habitats, we recommend that unburned patches of habitat -- refugia -- be spared during any one prescribed burn and that the rotation between burn units be long enough to allow for effective emigration/recolonization before the refugia are themselves burned.