



Platypolia anceps Gray-dusted Sallow



n=7
High Mt.

n=10
Low Mt.

n=0
Pd




High counts of:

- 3 - Swain - 2003-10-08
- 2 - Haywood - 2002-10-09
- 2 - Swain - 1988-10-11

● = Sighting or Collection in NC since 2005
● = Not seen since 2005

Status Rank
NC US NC Global



FAMILY: Noctuidae SUBFAMILY: Noctuinae TRIBE: Xylenini
TAXONOMIC_COMMENTS:

FIELD GUIDE DESCRIPTIONS:

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS:

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS:

DISTRIBUTION: Please refer to the dot map.

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT: Our records come from stands of northern hardwoods and cove forests.

FOOD: Larvae have been found on Bigleaf Aster (*Eurybia macrophylla*) and have been reared on blueberry and cherry (Wagner et al., 2011). In North Carolina, a larva has been recorded feeding on solomon's-seal (*Polygonatum*).

OBSERVATION_METHODS:

NATURAL HERITAGE PROGRAM RANKS: G5 SNR [S1S2]

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: