



# *Sigmoria areolata* No common name



High counts of:

- 6 - Buncombe - 1977-09-08
- 6 - Buncombe - 1977-09-08
- 6 - Buncombe - 1977-09-09

Status	Rank		
NC	US	NC	Global

n=0  
CP

FAMILY: Xystodesmidae SUBFAMILY: TRIBE:  
TAXONOMIC COMMENTS:

FIELD GUIDE DESCRIPTIONS:  
ONLINE PHOTOS:

ID COMMENTS:

DISTRIBUTION: Shelley (1981) regarded this species as having a very restricted distribution: "Sigmoria areolata is a highly localized endemic species known only from the area around Baldwin Gap in central Buncombe county. This area was revisited in June 1979, but no specimens were encountered"""

HABITAT: Shelley (1981) found the type specimens in hardwood forest growing on a hillside adjacent to a cove forest. He speculated that it occupies the same range of habitats as latior, which includes a range of dry-to-wet hardwood forests, from hillsides to floodplains. However, this species -- unlike areolata -- appears to have a much more restricted geographic range, which needs to be explained in terms of its habitat associations.

OBSERVATION METHODS:

NATURAL HERITAGE PROGRAM RANKS: [GNR] [SH]

STATE PROTECTION:

COMMENTS: This species appears to be endemic to a very small area and appears to be known only from historic records; Shelley (1981) reported he couldn't find the species when he searched for them at the type locality in 1979.