



Little Glassywing *Vernia verna*



Status and Rank	
State	Global
S5	G5

DISTRIBUTION: Essentially statewide, occurring in all three provinces, but very few records from the southeastern portion of the state. However, there are numerous records for coastal SC; thus, it should occur in most (if not all) Coastal Plain counties in NC, but it is certainly not an artifact of field work that there are so few records for parts of the Coastal Plain.

ABUNDANCE: Fairly common and widespread, to locally common, in the Mountains and Piedmont. Uncommon in the northern and western Coastal Plain (including the Sandhills); seemingly rare in the outer (eastern) Coastal Plain, and very rare to rare in parts of the central Coastal Plain. Declines as seen in many other grass skippers in recent years are not overly apparent with the Little Glassywing, but it likely is slightly less numerous than it was 20 or more years ago.

FLIGHT PERIOD: Two broods downstate; mid-May to early July, and mid- or late July to late September, rarely to mid-October. This species has a distinct gap in the flight period in early summer in the Piedmont and Coastal Plain. However, it flies in the Mountains in July; the primary brood there is from late May or early June to early August. There is a small second brood in the Mountains in late August and September, but more data needed for these months.

HABITAT: This is a species of mesic to somewhat damp places, and is not numerous in dry sites. It also has a tendency toward wooded edges and is not characteristic of open fields. It is usually seen along borders of hardwood or mixed forests, along powerline clearings through forests, along roads and wide trails through forests, and so forth. In the Coastal Plain, it favors damp places more so than in the upstate.

FOOD AND NECTAR PLANTS: Tall Redtop (*Tridens flavus*) is the most often reported foodplant, but other tall native grasses are likely used. The species nectars on a great variety of flowers; mountain-mints (*Pycnanthemum* spp.) are a favorite in the Piedmont, and Indian-hemp (*Apocynum cannabinum*) is used throughout the state.

COMMENTS: This species can be difficult to separate from both the Northern Broken-dash and the Dun Skipper; worn individuals can also be confused with the Crossline Skipper. Powerline clearings are good places to find them. The species can be surprisingly local; you might find 5-10 in an hour at a locale, but it can be hard to find in seemingly similar-looking places elsewhere. Nonetheless, it is a numerous skipper in NC.

It is difficult to explain the "gap" in records for the southern Coastal Plain, considering its presence in a great number of Coastal Plain counties in SC. This has not been for lack of field work in NC, and margins of moist hardwood forests, powerline cuts, etc., are abundant. Is Tall Redtop or other foodplants scarce there?