A REPTILE Checklist for Hanging Rock State Park 29 taxa

Herpetology is the branch of zoology dedicated to the study of amphibians (frogs, toads, salamanders, and newts) and of reptiles (snakes, lizards, turtles, and crocodilians). "Herps" are a diverse and interesting group of animals that are important elements in the fabric of life.

Amphibian is derived from the Greek work "amphibios". "Amphi" meaning 'both' and "bio" meaning 'life' - hence leading a double life. Indeed, most amphibians are aquatic as juveniles and breath with gills. After transformation, the adults usually live on land and most have lungs.

Many amphibians are very sensitive to environmental change. Their diversity and abundance can be an indication of an area's ecological health.

In early spring and during the summer listen for the calls of frogs and toads. Salamanders and newts, however, are silent and can be quiet secretive.

Reptiles, unlike amphibians, have dry skin covered with horny scales. Most, with the exception of snakes and legless lizards, have 4 legs with five clawed digits on each foot. While some amphibians retain gills as adults, all reptiles breathe solely by means of lungs.

Look for these cold-blooded animals on warm to hot sunny days when they bask on rocks, logs and tree trunks. Sixty-seven species have been recorded in North Carolina.

> For more information, contact: Hanging Rock State Park

(336) 593-8480 hanging.rock@ncparks.gov

Notes

REPTILE CHECKLIST Hanging Rock State Park



Seasons:

Spring - March to May Summer - June to August Fall -September to November Winter - December to February

Web Links

Amphibians and Reptiles of North Carolina State Parks https://auth1.dpr.ncparks.gov/nrid/public.php

Resources

Snakes of the Southeast by Whit Gibbons and Mike Dorcas Univ. of Georgia Press http://www.ugapress.org

Amphibians and Reptiles of the Carolinas and Virginia by Jeffrey C. Beane, Alvin L. Braswell, Joseph C. Mitchell, and William M. Palmer Univ. of North Carolina Press http://uncpress.unc.edu

Reptiles of North Carolina by William M. Palmer and Alvin L. Braswell Univ. of North Carolina Press http://uncpress.unc.edu



https://www.ncparks.gov/junior-rangers

| 29 REPTILE taxa | | | | |
|--|-----|-----|-----|---|
| EMYDIDAE: [2] | | | | |
| Eastern Box Turtle | х | х | х | - |
| Terrapene carolina | | | | |
| KINOSTERNIDAE: [1] | | | | |
| Common Musk Turtle (Stinkpot) | Х | - | - | - |
| Sternotherus odoratus | | | | |
| COLUBRIDAE: [16] | | | | |
| Central/Eastern Ratsnake Complex | х | х | х | - |
| Pantherophis alleghaniensis-quadrivittatus | | | | |
| Ring-necked Snake | х | х | х | - |
| Diadophis punctatus | | | | |
| Eastern Hog-nosed Snake | х | x | - | - |
| Heterodon platirhinos | | | | |
| Queen Snake | x | x | x | - |
| Regina septemvittata | | | | |
| Brown Snake | - | x | - | - |
| Storeria dekayi | | | | |
| Southeastern Crowned Snake | x | x | - | - |
| Tantilla coronata | | | | |
| VIPERIDAE: [2] | | | | |
| Copperhead | х | x | х | _ |
| Agkistrodon contortrix | | | | |
| Timber Rattlesnake | x | x | x | _ |
| Crotalus horridus | | | | |
| COLUBRIDAE: [16] | | | | |
| Smooth Earthsnake | x | _ | _ | _ |
| — Virginia valeriae | | | | |
| Eastern Worm Snake | x | x | x | _ |
| | | | | |
| Northern Scarlet Snake | _ | x | _ | _ |
| | | | | |
| Northern Black Racer | x | x | x | _ |
| Coluber constrictor | | | | |
| Scarlet Kingsnake | x | x | x | x |
| Lampropeltis elapsoides | | | | |
| Eastern Kingsnake | _ | _ | _ | _ |
| Lampropeltis getula | | | | |
| Northern Water Snake | v | v | _ | _ |
| Nerodia sipedon | Λ | Λ | | |
| Northern Rough Green Snake | v | v | _ | _ |
| Opheodrys aestivus | ^ | ^ | _ | - |
| Corn Snake | 3.5 | 3.5 | 3.5 | |
| | X | X | X | _ |
| Pantherophis guttatus | | | | |

| | S | S | F | И |
|---------------------------------------|---|---|---|---|
| COLUBRIDAE: [16] | | | | |
| Northern Red-bellied Snake | x | x | x | _ |
| Storeria occipitomaculata | | | | |
| Common Gartersnake | x | x | x | _ |
| Thamnophis sirtalis | | | | |
| PHRYNOSOMATIDAE: [1] | | | | |
| Eastern Fence Lizard | x | x | х | _ |
| Sceloporus undulatus | | | | |
| SCINCIDAE: [3] | | | | |
| Common Five-lined Skink | x | x | х | х |
| Plestiodon fasciatus | | | | |
| Southeastern Five-lined Skink | _ | _ | _ | _ |
| Plestiodon inexpectatus | | | | |
| Broad-headed Skink | x | x | _ | _ |
| Plestiodon laticeps | | | | |
| TEIIDAE: [1] | | | | |
| Eastern Six-lined Racerunner | x | _ | _ | _ |
| Aspidoscelis sexlineatus | | | | |
| CHELYDRIDAE: [1] | | | | |
| Eastern Snapping Turtle | x | x | _ | _ |
| Chelydra serpentina | | | | |
| EMYDIDAE: [2] | | | | |
| Painted Turtle | _ | x | _ | _ |
| Chrysemys picta | | | | |
| River Cooter | x | _ | _ | _ |
| Pseudemys concinna | | | | |
| , , , , , , , , , , , , , , , , , , , | | | | |