



Argia apicalis Blue-fronted Dancer



NC counties: 95 or 95%
Number of records added in 2023 = 45

● = Sighting or Vetted in NC since 2004
● = Not seen since 2004

High counts of:
175 - Chatham - 2022-07-11
100 - Chatham - 2010-07-11
100 - Wake - 2021-07-21

Earliest date: Union 2015-04-14
Latest date: Edgecombe 2017-10-28

Status		Rank	
NC	US	NC	Global
-	-	S5	G5

DISTRIBUTION: Essentially statewide, though possibly absent in the eastern Tidewater Region. Presumably absent on the Outer Banks.

ABUNDANCE: Often very common in the Piedmont and western Coastal Plain, particularly around flowing water in the Piedmont. Fairly common to locally common in the mountains and the central and southern Coastal Plain. Rare in the northeastern and far eastern Coastal Plain, and probably absent near the coast there.

FLIGHT: Early May through late October in the Piedmont and the Coastal Plain. The flight in the mountains is from mid-May at least to mid-September (and likely later).

HABITAT: Favors rivers and large streams, but can be found away from water. Also occurs around lake and pond margins.

BEHAVIOR: Prefers open areas with sunlight, using rocks, ground, or available vegetation for perching. It seems to be the most commonly seen dancer on dirt roads and other bare ground well away from water.

COMMENTS: This is one of our most common and widespread damselflies, except near the northeastern and eastern coastal areas. There are several one-day counts of at least 50 individuals, all from the lower Piedmont.