



Epitheca cynosura Common Baskettail



NC counties: 100 or 100%
Number of records added in 2023 = 69

● = Sighting or Vetted in NC since 2004
● = Not seen since 2004

High counts of:
400 - Camden - 2017-04-13
150 - Onslow - 2010-04-24
130 - Wake - 2013-05-03

Earliest date: Bladen 2023-03-04
Latest date: Macon 1981-08-05

Status		Rank	
NC	US	NC	Global
-	-	S5	G5

DISTRIBUTION: Statewide; occurs in every NC county, including those along the immediate coast.

ABUNDANCE: Common to occasionally abundant across the state. Often the most numerous species seen on an outing in the spring season. Dozens can sometimes be seen in a day.

FLIGHT: Spring/early summer season only; mid-March to late June or early July downstate, and to early August in the mountains (where the flight begins in late March). It is infrequent downstate after the end of May.

HABITAT: Mainly at still or slow-moving water of ponds, lakes, and pools; less so at slow creeks and rivers.

BEHAVIOR: Males are often seen flying low over pools and ponds, and at times over smaller creeks, perching quite frequently along the margins. Away from water, adults range widely to woodland margins, sunny roads and trails, and so forth, where they perch unwarily on twigs or other low vegetation and thus can be studied closely.

COMMENTS: Identification of most baskettails (except Prince) can be tricky, as some Mantleds, most Slender, and especially Robust baskettails closely resemble the Common Baskettail. In fact, there is probably no single field mark that might identify a dragonfly as a Common; a suite of marks, such as abdomen shape, abdomen width, and wing coloration must be used. Nonetheless, observers will quickly tire of seeing this species in spring, in hopes of finding less numerous baskettails, darners, and skimmers.